

SAFETY SEX

Each soft-gel capsule contains:

Composition:

Equivalent to Clindamycin	
Clotrimazole I.P	J
Metronidazole	1

Indications: Above 18 year's Female Vaginal use, Fungicidal, Bacterial, Trichmonacidal.

Description:

Clindamycin has a bacteriostatic effect. Clindamycin interferes with bacterial protein synthesis, in a similar way as erythromycin and chloramphenicol, by binding to the 50S subunit of the bacterial ribosome. This causes antagonism if administered simultaneously and possible cross-resistance.

Clotrimazole damages the fungal cell wall and alters the permeability. It also inhibits the activity of intracellular enzymes, leading to a buildup of toxic concentrations of hydrogen peroxide within the fungal cell, causing death of the cell. Systemic absorption is minimal.

Metronidazole is rather lipophilic and is rapidly and widely distributed after absorption. It is distributed to most body tissues and fluids, including to bone, abscesses, the CNS, and seminal fluid. It is less than 20% bound to plasma proteins in humans.

Side Effects

Local cutaneous effects may include irritation, hypersensitivity, burning, pruritus, erythema, fissuring, or swelling.

Dosage: As per the physician's advice.

Presentations: 3 soft gel Capsules

MRP	Retailer	Stockist
54.00	43.20	38.88

Taj Group of Companies

